

Перелік продукції ПрАТ «Фармацевтична фірма «Дарниця»

№	DRUG NAME	INN	PRESCRIPTION / OTC	ATC	Therapeutic indications
1	ADRENALIN-DARNITSA, solution for injection 1.8 mg/ml, 10 x 1 ml	Epinephrine	<i>Rx</i>	C01C A24	Non-glycoside cardiotoxic drug. Drug for emergency treatment. It is used in anaphylactic shock to relieve asthma attack, in asystole, cardiac arrest, acute 3rd degree AV block; prolongs the action of local anaesthetics.
2	AZICIN®, 6 x 250 mg caps.	Azithromycin	<i>Rx</i>	J01F A10	Macrolide antibiotic. It is used to treat respiratory tract infections, skin and soft tissue infections, and sexually transmitted infections. The drug is prescribed for adults and children weighing more than 45 kg.
3	AZICIN®, 3 x 500 mg film-coated tabs.	Azithromycin	<i>Rx</i>	J01F A10	Macrolide antibiotic. It is used to treat respiratory tract infections, skin and soft tissue infections, and sexually transmitted infections. The drug is prescribed for adults and children weighing more than 45 kg.
4	ALOE EXTRACT LIQUID - DARNITSA, 10 x 1 ml amps. for injection	Aloes	<i>Rx</i>	A16A X10	Biogenic stimulator. It has adaptogenic and general tonic effect. It is used in ophthalmology for the treatment of inflammatory eye diseases, and in the combination therapy of gastric ulcer and duodenal ulcer.
5	AMBROXOL-DARNITSA, 20 x 30 mg tabs.	Ambroxol	<i>OTC</i>	R05C B06	Mucolytic drug. It is used for secretolytic therapy of acute and chronic bronchopulmonary diseases in adults and children aged over 6 years.
6	AMLOPRIL®- DARNITSA, 20 x 5 mg tablets	Amlodipin	<i>Rx</i>	C08C A01	Calcium antagonist. Due to the vasodilator effect on peripheral arterioles and coronary vessels it causes hypotensive and antianginal effects. It is used in arterial hypertension, stable exertional angina, Prinzmetal angina.
7	AMLOPRIL®- DARNITSA, 20 x 10 mg tablets	Amlodipin	<i>Rx</i>	C08C A01	Calcium antagonist. Due to the vasodilator effect on peripheral arterioles and coronary vessels it causes hypotensive and antianginal effects. It is used in arterial hypertension, stable exertional angina, Prinzmetal angina.
8	ANALGIN-DARNITSA, solution for injection 500 mg/ml, 10 x 2 ml	Metamizole Sodium	<i>Rx</i>	N02B B02	Analgesic and antipyretic drug. It is used in pain syndrome, fever, renal and hepatic colic in combination with antispasmodics.
9	ANALGIN-DARNITSA, 10 x 500 mg tabs.	Metamizole Sodium	<i>OTC</i>	N02B B02	Analgesic and antipyretic drug. It is used to relieve pain and in fever. The duration of use is not more than 3 days. The drug is prescribed for adults and children aged over 12 years.

10	ANTISTRUMIN-DARNITSA, 100 x 1 mg tabs.	Potassium Iodide	<i>Rx</i>	H03C A	Iodine drug. The drug is prescribed for prevention and treatment of iodine deficiency disorders and thyroid diseases in adults and children aged over 12 years.
11	ANTISTRUMIN-DARNITSA, 50 x 1 mg tabs.	Potassium Iodide	<i>Rx</i>	H03C A	Iodine drug. The drug is prescribed for prevention and treatment of iodine deficiency disorders and thyroid diseases in adults and children aged over 12 years.
12	ARGININE-DARNITSA, solution for injection 40 mg/ml, 10 x 5 ml	Arginine Glutamate	<i>Rx</i>	A05B A01	The drug affects the metabolism in the liver cells, helping to neutralize and eliminate toxic metabolites and improving energy supply of hepatocytes. The drug is indicated for use in hepatic encephalopathy, in acute alcohol poisoning, in the combination therapy of acute and chronic hepatitis of different aetiology, liver cirrhosis. The drug is prescribed in complications (late gestosis, fetoplacental insufficiency, hepatobiliary pathology) in the third trimester of pregnancy.
13	ASCORBIC ACID-DARNITSA, solution for injection 100 mg/ml, 10 x 2 ml	Ascorbic Acid	<i>Rx</i>	A11G A01	Vitamin C. It is used in hypovitaminosis C, scurvy, bleedings, haemorrhagic diathesis, overdose with anticoagulants, infectious diseases, intoxications, dystrophy, nephropathy during pregnancy, Addisonian crisis, bone fractures and slow-healing wounds, in increased brain stress and heavy physical labour.
14	ASCORBIC ACID-DARNITSA, solution for injection 50 mg/ml, 10 x 2 ml	Ascorbic Acid	<i>Rx</i>	A11G A01	Vitamin C. It is used in hypovitaminosis C, scurvy, bleedings, haemorrhagic diathesis, overdose with anticoagulants, infectious diseases, intoxications, dystrophy, nephropathy during pregnancy, Addisonian crisis, bone fractures and slow-healing wounds, in increased brain stress and heavy physical labour.
15	ASCOPHENUM-DARNITSA, 6 tabs.	Paracetamol Caffeine Aspirin	<i>OTC</i>	N02B A51	Combination drug with analgesic, antipyretic and anti-inflammatory effect. It is used to treat mild or moderate pain syndrome, as well as hyperthermia.
16	ATROPINE-DARNITSA®, solution for injection 1 mg/ml, 10 x 1 ml	Atropine Sulfate	<i>Rx</i>	A03B A01	M-cholinoblocker. It reduces the secretion of exocrine glands and the tone of smooth muscles of internal organs. The drug is prescribed in stomach ulcers and duodenal ulcers, gastrointestinal spasms and urinary tract spasms, bronchial asthma, bradycardia. It is used before general anaesthesia and during surgery. It is a specific antidote in poisoning with cholinomimetic compounds and anticholinesterase (including organophosphate) substances.
17	ACETYLSALICYLIC ACID-DARNITSA, 10 x 500 mg tabs.	Acetylsalicylic Acid	<i>OTC</i>	N02B A01	Nonsteroidal anti-inflammatory drug. It is used to treat mild to moderate pain syndrome, feverish conditions, rheumatic diseases. In median doses it shows analgesic and antipyretic effects.

18	ACICLOVIR-DARNITSA, 20 x 200 mg tabs.	Aciclovir	<i>Rx</i>	J05A B01	Antiviral drug. It is used to treat infections caused by Herpes simplex virus and Varicella zoster virus and to prevent recurrence of infections caused by Herpes simplex in adults and children aged over 2 years.
19	BENZYL BENZOATE-DARNITSA, 250 mg/g ointment 30 g tube	Benzyl Benzoate	<i>OTC</i>	P03A X01	The drug acting against ectoparasites. It is used to treat scabies in adults and children aged over 3 years.
20	BETAMETHASONE-DARNITSA, cream 15 g tube	Betamethasone Valerate Tsetylpirydyinyu Chloride	<i>Rx</i>	D07B C01	Combined corticosteroid drug with antiseptic. The drug is prescribed in eczema, neurodermatitis, dermatitis, psoriasis (except widespread psoriasis), anogenital and senile pruritus.
21	BISACODYL-DARNITSA, 30 x 5 mg film-coated tabs.	Bisacodyl	<i>OTC</i>	A06A B02	Laxative. It is used for short-term (no more than 7-10 days) treatment of constipation in adults and children aged over 4 years.
22	BROMHEXINE-DARNITSA, 10 x 8 mg tabs.	Bromhexine	<i>OTC</i>	R05C B02	Mucolytic drug. It is used to facilitate expectoration in acute and chronic bronchopulmonary diseases in adults and children aged over 6 years.
23	BROMHEXINE-DARNITSA, 20 x 8 mg tabs.	Bromhexine	<i>OTC</i>	R05C B02	Mucolytic drug. It is used to facilitate expectoration in acute and chronic bronchopulmonary diseases in adults and children aged over 6 years.
24	BROMHEXINE-DARNITSA, 50 x 8 mg tabs.	Bromhexine	<i>OTC</i>	R05C B02	Mucolytic drug. It is used to facilitate expectoration in acute and chronic bronchopulmonary diseases in adults and children aged over 6 years.
25	VALIDOL-DARNITSA, 10 x 60 mg tabs.	A Solution Of Menthol In The Air Mentyllovomu Isovaleric Acid	<i>OTC</i>	C01E X	Combined cardiovascular drug. It is used in functional cardialgia of angioneurotic character, neurocirculatory dystonia of cardiac type, mild short-term anginal attacks, neurosis, hysteria, sea and air sickness, nausea of neurotic character.
26	VALIDOL-DARNITSA, 6 x 60 mg tabs.	A Solution Of Menthol In The Air Mentyllovomu Isovaleric Acid	<i>OTC</i>	C01E X	Combined cardiovascular drug. It is used in functional cardialgia of angioneurotic character, neurocirculatory dystonia of cardiac type, mild short-term anginal attacks, neurosis, hysteria, sea and air sickness, nausea of neurotic character.
27	VERAPAMIL-DARNITSA, 20 x 40 mg film-coated tabs.	Verapamil	<i>Rx</i>	C08D A01	Calcium antagonist. The drug is prescribed in angina pectoris (exertional, unstable, vasospastic, postinfarction), arrhythmia, arterial hypertension.
28	VERAPAMIL-DARNITSA, 50 x 80 mg film-coated tabs.	Verapamil	<i>Rx</i>	C08D A01	Calcium antagonist. The drug is prescribed in angina pectoris (exertional, unstable, vasospastic, postinfarction), arrhythmia, arterial hypertension.

29	VERAPAMIL HYDROCHLORIDE- DARNITSA, solution for injection 2.5 mg/ml, 10 x 2 ml	Verapamil	<i>Rx</i>	C08D A01	Calcium antagonist. It is used in paroxysmal supraventricular tachycardia, atrial flutter/atrial fibrillation (excluding WPW syndrome), unstable angina in patients with contraindications to nitrates and/or β -adrenoblockers, including vasospastic angina, variant angina, Prinzmetal angina, arterial hypertension, hypertensive crisis; in paediatric practice – in paroxysmal supraventricular tachycardia.
30	VICASOLUM- DARNITSA, solution for injection 10 mg/ml, 10 x 1 ml	Menadione Sodium Bisulfite	<i>Rx</i>	B02B A02	Vitamin K. It is used in bleeding associated with hypothrombinaemia, in bleeding in various pathologies, after surgeries, against overdose with phenylin, neodicumarinum, other anticoagulants - vitamin K antagonists.
31	VINPOCETINUM- DARNITSA, concentrate for solution for infusion 5 mg/ml, 10 x 2 ml	Vinpocetine	<i>Rx</i>	N06B X18	Psychostimulant and nootropic drug. It is used in cerebrovascular diseases, chronic vascular diseases of the choroid and retina, deafness, Meniere's disease.
32	VINPOCETINUM- DARNITSA, 30 x 5 mg tabs.	Vinpocetine	<i>Rx</i>	N06B X18	Psychostimulant and nootropic drug. It is used in cerebrovascular diseases, chronic vascular diseases of the choroid and retina, deafness, Meniere' disease.
33	WATER FOR INJECTIONS- DARNITSA, water for injections, 10 x 5 ml amps.	Water For Injection	<i>OTC</i>	V07A B	Solvent. It is used for the preparation of sterile solutions of medicinal and diagnostic products for subcutaneous, intramuscular or intravenous administration.
34	WATER FOR INJECTIONS- DARNITSA, water for injections, 10 x 2 ml amps.	Water For Injection	<i>OTC</i>	V07A B	Solvent. It is used for the preparation of sterile solutions of medicinal and diagnostic products for subcutaneous, intramuscular or intravenous administration.
35	GENTAMICIN SULFATE- DARNITSA, solution for injection 40 mg/ml, 10 x 2 ml	Gentamicin	<i>Rx</i>	J01G B03	Aminoglycoside antibiotic. It is used in bacterial infections caused by microflora sensitive to gentamicin, including: infections of the lower respiratory tract, complicated urogenital infections, bone and joint infections, skin and soft tissue infections, infected burn wounds, abdominal infection, infections of the central nervous system, septicaemia.
36	HEPARIN- DARNITSA, 600 IU/g gel 30 g tube	Heparin Sodium	<i>OTC</i>	C05B A53	Direct-acting anticoagulant. It is used for local treatment of varicose veins, thrombophlebitis of superficial veins, post-injection phlebitis, surface periphlebitis, complications after surgeries on veins, injuries of soft tissues, tendons and joints without damage of the skin integrity, subcutaneous haematomas.
37	GLUCOSE- DARNITSA, solution 400 mg/ml, 10 x 20 ml	Dextrose	<i>Rx</i>	B05B A03	Carbohydrate supply drug. It is used in hypoglycaemia.

38	GLUCOSE-DARNITSA, solution for infusion 50 mg/ml, 1 x 200 ml vial	Dextrose	<i>Rx</i>	B05C X01	The drug is indicated for use in intoxications, hypoglycaemia, hyper- and isotonic dehydration in children during surgeries to prevent water-electrolyte imbalance. The drug is used as a solvent of other compatible drug solutions.
39	GLUCOSE-DARNITSA, solution for infusion 50 mg/ml, 1 x 400 ml vial	Dextrose	<i>Rx</i>	B05C X01	The drug is indicated for use in intoxications, hypoglycaemia, hyper- and isotonic dehydration in children during surgeries to prevent water-electrolyte imbalance. The drug is used as a solvent of other compatible drug solutions.
40	DARSIL-DARNITSA, 100 x 22.5 mg film-coated tabs.	Silimarin	<i>OTC</i>	A05B A03	Hepatotrophic drug. It is used in chronic toxic liver injuries, as a part of a combined treatment of patients with chronic inflammatory liver disease or liver cirrhosis. The drug is prescribed for adults and children aged over 5 years.
41	DARSIL-DARNITSA, 50 x 22.5 mg film-coated tabs.	Silimarin	<i>OTC</i>	A05B A03	Hepatotrophic drug. It is used in chronic toxic liver injuries, as a part of a combined treatment of patients with chronic inflammatory liver disease or liver cirrhosis. The drug is prescribed for adults and children aged over 5 years.
42	DARSIL-DARNITSA, 30 x 22.5 mg film-coated tabs.	Silimarin	<i>OTC</i>	A05B A03	Hepatotrophic drug. It is used in chronic toxic liver injuries, as part of a combined treatment of patients with chronic inflammatory liver disease or liver cirrhosis. The drug is prescribed for adults and children aged over 5 years.
43	DEXAMETHASONE-DARNITSA, 1 mg/ml eye drops, 10 ml	Dexamethasone	<i>Rx</i>	S01B A01	Synthetic glucocorticosteroid. It is used in non-infectious inflammatory and allergic conditions of the conjunctiva, cornea and anterior segment of the eye, including inflammation reactions in the postoperative period.
44	DEXAMETHASONE-DARNITSA, solution for injection 4 mg/ml, 5 x 1 ml	Dexamethasone	<i>Rx</i>	H02A B02	Synthetic glucocorticosteroid. It is used in a number of endocrine diseases, in rheumatic, allergic diseases, skin diseases, diseases of gastrointestinal tract, respiratory system, kidneys, eyes, in oncological and haematological diseases.
45	DEXAMETHASONE-DARNITSA, 50 x 0.5 mg tabs.	Dexamethasone	<i>Rx</i>	H02A B02	Synthetic glucocorticosteroid. It is used in a number of endocrine diseases, in rheumatic, allergic diseases, skin diseases, diseases of gastrointestinal tract, respiratory system, nervous system, kidneys, eyes, in oncological and haematological diseases.

46	DIBAZOL-DARNITSA, solution for injection 10 mg/ml, 10 x 1 ml	Bendazol	<i>Rx</i>	C04A X31	Peripheral vasodilator. It is used in spasms of blood vessels (exacerbation of hypertensive disease, hypertensive crisis), and smooth muscles of internal organs (pyloric and intestinal spasms), in treatment of nervous system diseases in adults and children aged over 12 years.
47	DIBAZOL-DARNITSA, solution for injection 10 mg/ml, 10 x 5 ml	Bendazol	<i>Rx</i>	C04A X31	Peripheral vasodilator. It is used in spasms of blood vessels (exacerbation of hypertensive disease, hypertensive crisis), and smooth muscles of internal organs (pyloric and intestinal spasms), in treatment of nervous system diseases in adults and children aged over 12 years.
48	DIBAZOL-DARNITSA, 10 x 20 mg tabs.	Bendazol	<i>OTC</i>	C04A X31	Peripheral vasodilator. It is used for arterial hypertension, nervous system diseases (residual effects of polio, peripheral facial nerve palsy, polyneuritis, flaccid paralysis syndrome).
49	DICLOFENAC-DARNITSA, solution for injection 25 mg/ml, 10 x 3 ml	Diclofenac	<i>Rx</i>	M01A B05	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used in rheumatism and rheumatic diseases, acute gout attacks, renal and biliary colic, pain and swelling after surgeries, in severe migraine attacks.
50	DICLOFENAC-DARNITSA, solution for injection 25 mg/ml, 5 x 3 ml	Diclofenac	<i>Rx</i>	M01A B05	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used in rheumatism and rheumatic diseases, acute gout attacks, renal and biliary colic, pain and swelling after surgeries, in severe migraine attacks.
51	DICLOFENAC-DARNITSA, 30 x 25 mg film-coated tabs.	Diclofenac	<i>Rx</i>	M01A B05	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used in inflammatory and degenerative forms of rheumatic diseases, in acute gout attacks, spinal pain syndrome, pain and swelling after surgeries, in gynaecological and ENT diseases with pain syndrome.
52	DIMEDROLUM-DARNITSA, solution for injection 10 mg/ml, 10 x 1 ml	Diphenhydramine	<i>Rx</i>	R06A A02	Antihistamine drug. It is used in allergic diseases, hypersensitivity reactions, including anaphylactic reactions, itching dermatoses, chorea, Meniere's disease, postoperative vomiting.
53	DIMEDROLUM-DARNITSA, 10 x 50 mg tabs.	Diphenhydramine	<i>Rx</i>	R06A A02	Antihistamine drug. It is used to treat allergic reactions, allergic eye diseases, for prevention and treatment of allergic complications under drug administration, blood transfusions. The drug is prescribed in chorea, movement disorders, Meniere's disease. It is used as a sedative and hypnotic agent in neurosis, neurasthenia, insomnia.

54	DINORIK®-DARNITSA, 10 film-coated tabs.	Atenolol Chlorthalidone	<i>Rx</i>	C07C B03	Combined drug of beta-blockers with diuretics. It is used in arterial hypertension.
55	DITHYLINUM-DARNITSA, solution for injection 20 mg/ml, 10 x 5 ml	Suxamethonium	<i>Rx</i>	M03A B01	Muscle relaxant. It is used for intubation of trachea, endoscopic procedures, short-term surgeries, requiring muscle relaxation (reduction of dislocations, reposition the bone fragments), to maintain myoplegia during surgeries (gynaecologic, thoracic, abdominal), elimination of the cramps in tetanus.
56	DIAZOLIN-DARNITSA, 10 x 100 mg tabs.	Mebhydrolin	<i>OTC</i>	R06A X15	Antihistamine drug. It is used to prevent and treat seasonal and allergic rhinitis, hay fever, urticaria, food and drug allergies, itchy dermatitis in adults and children aged over 5 years.
57	DIOXISOL®-DARNITSA, solution, 100 g vial	Dioksydyn Lidocaine Hydrochloride	<i>OTC</i>	D03A X50	Combined drug with antibacterial action. It is used to treat infected wounds of soft tissues, 2nd to 4th degree burns, osteomyelitis. It is used in preparing wounds for autodermoplasty, for prevention of purulent-inflammatory processes in surgery and combustiology when performing plastic surgeries.
58	DIOXISOL®-DARNITSA, solution, 50 g vial	Dioksydyn Lidocaine Hydrochloride	<i>OTC</i>	D03A X50	Combined drug with antibacterial action. It is used to treat infected wounds of soft tissues, 2nd to 4th degree burns, osteomyelitis. It is used in preparing wounds for autodermoplasty, for prevention of purulent-inflammatory processes in surgery and combustiology when performing plastic surgeries.
59	DOXYCYCLINE-DARNITSA, 10 x 100 mg caps.	Doxycycline	<i>Rx</i>	J01A A02	Tetracycline antibiotic. It is used to treat infections caused by microorganisms susceptible to doxycycline in adults and children aged over 12 years.
60	DOFAMIN-DARNITSA, concentrate for solution for infusion 5 mg/ml, 10 x 5 ml	Dopamine	<i>Rx</i>	C01C A04	Non-glycoside cardiotoxic drug. It is used in acute hypotension, shock, decompensated chronic heart failure.
61	DOFAMIN-DARNITSA, concentrate for solution for infusion 40 mg/ml, 10 x 5 ml	Dopamine	<i>Rx</i>	C01C A04	Non-glycoside cardiotoxic drug. It is used in acute hypotension, shock, decompensated chronic heart failure.
62	DROTAVERINE-DARNITSA, solution for injection 20 mg/ml, 5 x 2 ml	Drotaverine	<i>Rx</i>	A03A D02	Myotropic antispasmodic. It is used in spasms of smooth muscles of the gastrointestinal tract, biliary system, urinary tract, in dysmenorrhoea.

63	DROTAVERINE-DARNITSA, 20 x 40 mg tabs.	Drotaverine	<i>OTC</i>	A03A D02	Myotropic antispasmodic. It is used in spasms of smooth muscles of the gastrointestinal tract, biliary system, urinary tract, in dysmenorrhoea, tension headache. It is prescribed for adults and children aged over 6 years.
64	DROTAVERINE-DARNITSA, 30 x 40 mg tabs.	Drotaverine	<i>OTC</i>	A03A D03	
65	ENALAPRIL-DARNITSA, 20 x 10 mg tabs.	Enalapril	<i>Rx</i>	C09A A02	ACE inhibitor. It is used for the treatment of essential and renovascular hypertension, heart failure in adults and children aged over 6 years. In patients with left ventricular dysfunction the drug helps to prevent the development of severe heart failure and coronary ischemia.
66	ENERGOTON®-DARNITSA, 60 x 20 mg film-coated tabs.	Trimetazidin	<i>Rx</i>	C01E B15	Cardiovascular drug. Optimizes ischemic cell energy metabolism. It is used for the symptomatic treatment of stable angina provided insufficient efficacy or intolerance to first-line antianginal drugs.
67	ETAMSYLATE-DARNITSA, solution for injection 125 mg/ml, 10 x 2 ml	Etamsylate	<i>Rx</i>	B02B X01	Antihemorrhagic drug. It is used for prevention and treatment of bleeding in adults and children.
68	EUPHYLLINE-DARNITSA, solution for injection 20 mg/ml, 10 x 5 ml	Theophylline	<i>Rx</i>	R03D A05	Bronchodilator, antispasmodic, vasodilator. It is used for bronchial obstruction syndrome in bronchial asthma, bronchitis, lung emphysema, disorders of the respiratory centre (night paroxysmal apnoea), in pulmonary heart. The drug is prescribed for adults and children aged over 14 years.
69	IBUPROFEN-DARNITSA, 50 x 200 mg tabs.	Ibuprofen	<i>OTC</i>	M01A E01	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used in the headache, toothache, migraine, dysmenorrhoea, neuralgia, back pain, joint pain, muscle pain, rheumatic pains, in feverish conditions, cold and flu symptoms. The drug is prescribed for adults and children aged over 12 years.
70	ISONIAZID-DARNITSA, solution for injection 100 mg/ml, 10 x 5 ml	Isoniazid	<i>Rx</i>	J04A C01	Antituberculosis drug. First-line drug for the treatment of all forms and localizations of tuberculosis in adults and children.

71	ISONIAZID-DARNITSA, 50 x 300 mg tabs.	Isoniazid	<i>Rx</i>	J04A C01	Antituberculosis drug. First-line drug for the treatment of all forms and localizations of tuberculosis (in combination with 3-4 other antituberculosis drugs) in adults and children aged over 4 years. The drug is prescribed as monotherapy for the treatment of latent tuberculosis infection and tuberculosis prevention in contact persons.
72	IMUSTAT, 10 x 100 mg film-coated tabs.	Umifenovirum	<i>OTC</i>	J05A X10	Antiviral drug. It is used for prevention and treatment of influenza A and B, acute respiratory infections, for combined therapy of chronic bronchitis, pneumonia and recurrent herpes infections in adults and children aged over 2 years; in combined therapy of acute intestinal infection of rotaviral aetiology in children.
73	CALCIUM GLUCONATE-DARNITSA (STABILIZED), solution for injection 100 mg/ml, 10 x 5 ml	Calcium Gluconate Calcium Levulinat	<i>Rx</i>	A12A A03	Calcium supplement. It is used in parathyroid gland failure, allergies, skin diseases, to decrease vascular permeability in various pathological processes, in liver diseases, hyperkalaemia, as a haemostatic agent and antidote for magnesium salt toxicity, toxicity of oxalic acid or its soluble salts, soluble salts of fluorine acid.
74	CALCIUM GLUCONATE-DARNITSA (STABILIZED), solution for injection 100 mg/ml, 10 x 10 ml	Calcium Gluconate Calcium Levulinat	<i>Rx</i>	A12A A03	Calcium supplement. It is used in parathyroid gland failure, allergies, skin diseases, to decrease vascular permeability in various pathological processes, in liver diseases, hyperkalaemia, as a haemostatic agent and antidote for magnesium salt toxicity, toxicity of oxalic acid or its soluble salts, soluble salts of fluorine acid.
75	CALCIUM GLUCONATE-DARNITSA, 10 x 500 mg tabs.	Calcium Gluconate	<i>OTC</i>	A12A A03	Calcium supplement. It is used in diseases accompanied by hypokalaemia, increased permeability of cell membranes, disturbance of nerve impulses in the muscle tissue, in the combined treatment of bleedings, in increased need for calcium, as an antidote for magnesium salt toxicity, toxicity of oxalic acid, soluble salts of fluorine acid.
76	CALCIUM CHLORIDE-DARNITSA, solution for injection 100 mg/ml, 10 x 5 ml	Calcium Chloride	<i>Rx</i>	B05X A07	Calcium supplement. It is used in cases of hypocalcaemia requiring rapid increase in the concentration of calcium ions in the blood plasma, in acute lead colic, in magnesium intoxication, hyperkalaemia.
77	CALCIUM CHLORIDE-DARNITSA, solution for injection 100 mg/ml, 10 x 10 ml	Calcium Chloride	<i>Rx</i>	B05X A07	Calcium supplement. It is used in cases of hypocalcaemia requiring rapid increase in the concentration of calcium ions in the blood plasma, in acute lead colic, in magnesium intoxication, hyperkalaemia.
78	CAPTOPRES 12.5-DARNITSA, 20 tabs.	Hydrochlorothiazide Captopril	<i>Rx</i>	C09B A01	Combined ACE inhibitor with diuretics. It is used in arterial hypertension.
79	CAPTOPRES-DARNITSA, 20 tabs.	Hydrochlorothiazide Captopril	<i>Rx</i>	C09B A01	Combined ACE inhibitor with diuretics. It is used in arterial hypertension.

80	CARBAMAZEPINE-DARNITSA, 20 x 200 mg tabs.	Carbamazepine	<i>Rx</i>	N03A F01	Antiepileptic drug. It is used for psychomotor epilepsy, acute mania, bipolar affective disorder, essential and multiple sclerosis, trigeminal neuralgia, essential glossopharyngeal neuralgia, diabetic neuropathy with pain syndrome, abstinence syndrome in patients with chronic alcoholism, diabetes insipidus of central origin, polydipsia and polyuria of neurohormonal origin.
81	CARBAMAZEPINE-DARNITSA, 50 x 200 mg tabs.	Carbamazepine	<i>Rx</i>	N03A F01	
82	CARBODERM-DARNITSA®, 100 mg cream 30 g tube	Urea	<i>OTC</i>	D08A X10	The drug has keratolytic, moisturizing, anti-itching, antibacterial and antifungal effects. It is used to treat skin diseases accompanied by increased formation of keratinizing epithelium, for treatment of psoriasis, seborrhoea, keratomycosis, hyperkeratosis, hyperkeratotic forms of eczema, ichthyosis.
83	KETozORAL®-DARNITSA, 2 mg/g shampoo 60 g vial	Ketoconazol	<i>OTC</i>	D01A C08	Antifungal drug. It is used to treat and prevent skin and hair lesions caused by Malassezia yeast microorganisms, such as dandruff, local pityriasis versicolor, seborrheic dermatitis.
84	KETOLONG-DARNITSA®, solution for injection 30 mg/ml, 10 x 1 ml	Ketorolac	<i>Rx</i>	M01A B15	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used to relieve moderate and severe postoperative pain during short time.
85	KETOLONG-DARNITSA®, 10 x 10 mg tabs.	Ketorolac	<i>Rx</i>	M01A B15	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used for short-term treatment of moderate pain, including postoperative pain.
86	ACIDUM AMINOCAPRONICUM-DARNITSA, solution for infusion 50 mg/ml, 1 x 100 ml vial	Acidum Aminocapronicum	<i>Rx</i>	B02A A01	Antihemorrhagic drug. It is used in surgeries and pathological processes accompanied by increased fibrinolytic activity of blood and tissue, to prevent secondary hypofibrinogenemia during massive transfusions of preserved blood.
87	CLARITHROMYCIN®, 14 x 500 mg film-coated tabs.	Clarithromycin	<i>Rx</i>	J01F A09	Macrolide antibiotic. It is used to treat infections caused by microorganisms susceptible to clarithromycin (respiratory tract infections, skin and soft tissue infections, odontogenic infections, mycobacterial infections, for eradication of H. pylori in patients with duodenal ulcer in the inhibition of secretion of hydrochloric acid). The drug is prescribed for adults and children aged over 12 years.

88	CLOPHELINUM-DARNITSA, 50 x 0.15 mg tabs.	Clonidine	<i>Rx</i>	C02A C01	Antihypertensive drug. It is used in hypertensive crisis (excluding hypertensive crisis with pheochromocytoma); rare for treatment of arterial hypertension (in combined therapy); in opioid withdrawal syndrome (in combined therapy).
89	KORVALOL-DARNYTSA®, oral drops, solution 25 ml x 1, bottle	Ethyl Ester And Acid-Bromizovaleria novoy, Phenobarbital, Oliya Peppermint	<i>OTC</i>	N05C B02	
90	KORVALOL-DARNYTSA®, oral drops, solution 25 ml x 1, bottle	Ethyl Ester And Acid-Bromizovaleria novoy, Phenobarbital, Oliya Peppermint	<i>Rx</i>	N05C B02	
91	CORDIAMIN-DARNITSA®, solution for injection 250 mg/ml, 10 x 2 ml	Nikethamide	<i>Rx</i>	R07A B02	Respiratory analeptic. It is used in collapse and asphyxia, in states of shock, acute and chronic disorders of blood circulation, in reduced vascular tone and respiratory depression in patients with infectious diseases and during recovery period, hypnotic toxicity and analgesic toxicity.
92	CAFFEINE-BENZOATE SODIUM-DARNITSA, solution for injection 100 mg/ml, 1 x 10 ml	Caffeine-Sodium Benzoate	<i>Rx</i>	N06B C01	Psycho-stimulant and nootropic drug. It is used in diseases accompanied by inhibition of the central nervous and cardiovascular systems, in respiratory depression, asphyxia, poisoning with drugs and other substances that inhibit the central nervous system, in asthenic syndrome, cerebral vasospasm.
93	LEVOMICETIN-DARNITSA, 10 x 250 mg tabs.	Chloramphenic ole	<i>Rx</i>	J01B A01	Broad-spectrum antibiotic. It is used to treat serious anaerobic infections (brain abscess) and abdominal infections; meningococcal infection, chlamydiosis, brucellosis, rickettsial infections such as typhus, ehrlichiosis, trachoma and Rocky Mountain spotted fever. It is also used in serious gastrointestinal infections, typhoid, paratyphoid, purulent wound infections, inguinal lymphogranuloma, severe infections caused by Haemophilus influenza, tularaemia, purulent otitis. It is prescribed for adults and children aged over 3 years.
94	LEVOMICETIN-DARNITSA, 10 x 500 mg tabs.	Chloramphenic ole	<i>Rx</i>	J01B A01	Broad-spectrum antibiotic. It is used to treat serious anaerobic infections (brain abscess) and abdominal infections; meningococcal infection, chlamydiosis, brucellosis, rickettsial infections such as typhus, ehrlichiosis, trachoma and Rocky Mountain spotted fever. It is also used in serious gastrointestinal infections, typhoid, paratyphoid, purulent wound infections, inguinal lymphogranuloma, severe infections caused by Haemophilus influenza, tularaemia, purulent otitis. It is prescribed for adults and children aged over 3 years.

95	LEFLOCK, 5 x 500 mg tabs.	Levofloxacin	<i>Rx</i>	J01M A12	Fluoroquinolone antibiotic. The drug is prescribed for adults to treat mild to moderate infectious and inflammatory diseases caused by microorganisms susceptible to the drug: Acute sinusitis, exacerbation of chronic bronchitis, infections of skin and soft tissue, in combined therapy of treatment-resistant forms of tuberculosis, community-acquired pneumonia, infections of kidneys and urinary tract infections, including pyelonephritis and prostatitis.
96	LEFLOCK-DARNITSA, solution for infusion 5 mg/ml, 1 x 100 ml vial	Levofloxacin	<i>Rx</i>	J01M A12	Fluoroquinolone antibiotic. The drug is prescribed for adults to treat inflammation in adults caused by bacteria susceptible to levofloxacin: pneumonia, complicated urinary tract infections, infections of skin and soft tissue, chronic bacterial prostatitis.
97	LIDOCAINE-DARNITSA, solution for injection 20 mg/ml, 10 x 2 ml	Lidocaine	<i>Rx</i>	N01B B02	Local anaesthetic. It is used for local anaesthesia (terminal, infiltration, conduction) in surgery, ophthalmology, dentistry, otolaryngology, for blockade of peripheral nerves and nerve plexus in various pain syndromes. The drug is prescribed for adults and children aged over 12 years.
98	LINCOMYCIN-DARNITSA, solution for injection 300 mg/ml, 10 x 1 ml	Lincomycin	<i>Rx</i>	J01F F02	Lincosamide antibiotic. It is used to treat adults and children aged over 1 month of infections caused by susceptible strains of gram-positive aerobic or anaerobic bacteria, including upper respiratory tract infections, respiratory tract infections, skin and soft tissue infections, infections of bones and joints, including osteomyelitis and septic arthritis, septicaemia and endocarditis, bacterial dysentery.
99	LINCOMYCIN-DARNITSA, solution for injection 300 mg/ml, 10 x 2 ml	Lincomycin	<i>Rx</i>	J01F F02	
100	LORATADINE-DARNITSA, 10 x 10 mg tabs.	Loratadine	<i>OTC</i>	R06A X13	Antihistamine drug. It is used for the symptomatic treatment of allergic rhinitis and chronic idiopathic urticaria in adults and children aged over 2 years.
101	MAGNESIUM SULFATE-DARNITSA, solution for injection 250 mg/ml, 10 x 10 ml	Magnesium Sulfate	<i>Rx</i>	B05X A05	Magnesium supplement. The drug has hypotensive, arterioldilating, antiarrhythmic, sedative, anticonvulsant, diuretic, antispasmodic, tocolytic effect. It is used in hypertensive crisis, ventricular heart rhythm disturbances (pirouette-type tachycardia), convulsive syndrome, eclampsia, hypomagnesaemia, in combined therapy of preterm labour, exertional angina, as antidote for poisoning with salts of heavy metals, tetraethyl lead, soluble barium salts.
102	MAGNESIUM SULFATE-DARNITSA, solution for injection 250 mg/ml, 5 x 10 ml	Magnesium Sulfate	<i>Rx</i>	B05X A05	

103	MEDICHRONAL®-DARNITSA, granules for oral use, compl. No.1	Glycine Glucose Monohydrate Sodium Formate	<i>OTC</i>	N07B B	The drug for prevention and treatment of alcohol intoxication. It is used for prevention of alcohol intoxication, reduces symptoms of alcohol intoxication, increases the functional activity of the cerebral cortex, has positive effect on liver function, improves general well-being.
104	MEDICHRONAL®-DARNITSA, granules for oral use, compl. No.21	Glycine Glucose Monohydrate Sodium Formate	<i>OTC</i>	N07B B	
105	MEDICHRONAL®-DARNITSA, granules for oral use, compl. No.7	Glycine Glucose Monohydrate Sodium Formate	<i>OTC</i>	N07B B	
106	METAMAX, 40 x 250 mg caps.	Meldonium	<i>Rx</i>	C05C X	Cardiovascular drug. It has cardioprotective, antianginal, antihypoxic effects. The drug is used in the combined therapy of coronary heart disease, cardialgia on the background dishormonal cardiomyopathy, cerebrovascular disturbances, decreased performance, physical overload, including in athletes, withdrawal syndrome in chronic alcoholism, acute disorders of blood circulation in the retina.
107	METAMAX, solution for injection 100 mg/ml, 10 x 5 ml	Meldonium	<i>Rx</i>	C05C X	Cardiovascular drug. It has cardioprotective, antianginal, antihypoxic effects. The drug is used in the combined therapy of ischemic heart disease, acute and chronic cerebrovascular, physical overload, abstinence syndrome in chronic alcoholism, haemophthalmia, haemorrhages in the retina of different aetiology, thrombosis of retinal central vein and its branches, retinopathies of different aetiology (diabetic, hypertensive).
108	METHYLURACIL & MYRAMISTIN, ointment, 15 g tube	Methyluracilum Myramistin	<i>OTC</i>	D03A X50	Drug for treatment of wounds and ulcers. It has antibacterial and wound-healing properties. It is used in complications, poor healing wounds, radiation and trophic ulcers (including patients with diabetic angiopathy, ulcers in debilitated patients with significant depression of immunity), burns and other skin injuries. The drug has sunscreen effect in patients with photodermatoses.
109	METHYLURACIL & MYRAMISTIN, ointment, 30 g tube	Methyluracilum Myramistin	<i>OTC</i>	D03A X50	
110	METOCLOPRAMIDE -DARNITSA, solution for injection 5 mg/ml, 10 x 2 ml	Metoclopramide	<i>Rx</i>	A03F A01	Peristaltic stimulator. Antiemetic action and the effect of accelerating gastric emptying and passing through the small intestine. The drug is used in dysmotility of upper gastrointestinal tract, nausea, urge to vomiting and vomiting (in migraine, liver and kidney diseases, traumatic brain injuries, drug

					intolerance), gastroparesis of diabetic genesis, to facilitate catheterization of duodenum and small intestine, to facilitate the emptying of the stomach and small intestine in X-ray examination of the stomach and small intestine.
111	METOCLOPRAMIDE-DARNITSA, 50 x 10 mg tabs.	Metoclopramide	<i>Rx</i>	A03F A01	Peristaltic stimulator. Antiemetic action and the effect of accelerating gastric emptying and passing through the small intestine. It is used as a second-line drug for the symptomatic treatment of nausea and vomiting, prevention of nausea and vomiting caused by radiotherapy, delayed (not acute) nausea and vomiting caused by chemotherapy, postoperative nausea and vomiting. It is prescribed for adults and children aged over 2 years.
112	METRONIDAZOLE-DARNITSA, solution for infusion 5 mg/ml, 1 x 100 ml vial	Metronidazole	<i>Rx</i>	J01X D01	Antibacterial drug. It is used to treat and prevent infections caused by microorganisms susceptible to metronidazole in adults and children aged over 2 years :infections of the central nervous system, lungs and pleura, gastrointestinal tract and abdominal infections, gynaecological infections, upper respiratory tract and oral cavity infections, infections of bones and joints, gas gangrene, septicaemia with thrombophlebitis; Prophylactic use is always indicated before surgeries with a high risk of anaerobic infections (before intra-abdominal and gynaecological surgeries).
113	MEFENAMIC ACID-DARNITSA, 20 x 500 mg tabs.	Mefenamic Acid	<i>OTC</i>	M01A G01	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used in acute upper respiratory infections and influenza, mild to moderate pain, in primary dysmenorrhoea, dysfunctional menorrhagia, including those caused by the presence of intrauterine contraceptive, in the absence of pelvic pathology, inflammatory diseases of the musculoskeletal system. It is prescribed for adults and children aged over 5 years.
114	MEFENAMYNKA, 20 x 500 mg film-coated tabs.	Mefenamic Acid	<i>OTC</i>	M01A G01	Nonsteroidal anti-inflammatory and anti-rheumatic agent. It is used in adults and children aged over 5 years with acute respiratory infections and influenza; for the relief of mild to moderate pain (muscular, articular, traumatic, dental, postoperative and postpartum, headaches of different aetiology); in primary dysmenorrhoea; dysfunctional menorrhagia without pelvic pathology; in inflammatory diseases of the musculoskeletal system.
115	MICONAZOLE-DARNITSA, 20 mg cream 15 g tube	Miconazole	<i>OTC</i>	D01A C02	Antifungal drug. It is for fungal infections of the skin and nails caused by dermatophytes or Candida fungi susceptible to miconazole, in superinfections caused by gram-positive bacteria.

116	MIORITMIL-DARNITSA, 30 x 200 mg tabs.	Amiodarone	<i>Rx</i>	C01B D01	Class III antiarrhythmic drug. It is used for prevention of relapse: - life-threatening ventricular tachycardia: the drug is prescribed in hospital environment under supervision; - reported symptomatic tachycardia and disabling ventricular tachycardia; - reported supraventricular tachycardia, if the need for prescription of the drug is identified in patients with disease resistant to other treatments, or in the presence of contraindications to other therapies of ventricular fibrillation. The drug is also prescribed for treatment of supraventricular tachycardia: slowdown or reduction of atrial fibrillation and atrial flutter.
117	MYRAMISTIN, 0.01 % solution, 50 ml	Myramistin	<i>OTC</i>	D08A J10	Antiseptic and disinfectant. It has a strong antimicrobial effect on gram-positive and gram-negative, aerobic and anaerobic, spore-forming and asporogenic bacteria, including hospital strains. The drug is detrimental to pathogens of sexually transmitted infections: gonococcus, treponema pallidum, trichomonas, chlamydia and herpes viruses, human immunodeficiency. It has antifungal action on a number of pathogenic fungi. Resistance of microorganisms to antibiotics is reduced under the influence of myramistin. It is used in surgery, traumatology, combustiology, obstetrics and gynaecology, dermatology, venereology, otorhinolaryngology, dentistry.
118	MYRAMISTIN®-DARNITSA, 5 mg ointment 1000 g jar	Myramistin	<i>OTC</i>	D08A J10	Antiseptic and disinfectant. The drug is used: -in surgery, traumatology: local treatment of infected wounds of different localization and aetiology, prevention of secondary infection of granulating wounds; -in combustiology: treatment of 2nd and 3rd-degree burns; preparation of burn wounds to dermoplasty;
119	MYRAMISTIN®-DARNITSA, 5 mg ointment 30 g tube	Myramistin	<i>OTC</i>	D08A J10	-in dermatology for treatment of strepto-, staphylo-derma, dermatomycosis of smooth skin, feet and large folds (including dyshidrotic forms and forms complicated by pyoderma), nails (onychomycosis), keratomycosis (including pityriasis versicolor), candidiasis of the skin and mucous membranes;
120	MYRAMISTIN®-DARNITSA, 5 mg ointment 15 g tube	Myramistin	<i>OTC</i>	D08A J10	-to prevent complications of wound infection in non-major industrial and home injuries.
121	SODIUM ADENOSINE TRIPHOSPHATE-DARNITSA, solution for injection 10mg/ml, 10 x 1 ml	Adenosine Triphosphate Disodium Salt	<i>Rx</i>	C01E B10	Cardiovascular drug. The drug has metabolic, antiarrhythmic effect, improves cerebral and coronary circulation.

122	SODIUM THIOSULFATE-DARNITSA, solution for injection 300mg/ml, 10 x 5 ml	Sodium Thiosulphate	<i>Rx</i>	V03A B06	Antidote. It has anti-toxic, anti-inflammatory and desensitizing effect. It is used poisoning with compounds of arsenic, mercury, lead, hydrocyanic acid and its salts, iodine, bromine and their salts.
123	SODIUM CHLORIDE-DARNITSA, solution for injection 9mg/ml, 10 x 10 ml	Sodium Chloride	<i>Rx</i>	B05X A03	It is used to supplement the lack of fluid in the body in a complex of intensive care measures, as a solvent of other compatible drugs. It can be applied topically for lavage of the wounds, nasal mucosa, as well as catheters and systems for transfusion.
124	SODIUM CHLORIDE-DARNITSA, solution for injection 9mg/ml, 10 x 5 ml	Sodium Chloride	<i>Rx</i>	B05X A03	
125	SODIUM CHLORIDE-DARNITSA, solution for injection 9mg/ml, 1 x 200 ml vial	Sodium Chloride	<i>Rx</i>	B05X A03	
126	SODIUM CHLORIDE-DARNITSA, solution for injection 9mg/ml, 1 x 400 ml vial	Sodium Chloride	<i>Rx</i>	B05X A03	Solution for infusion. It is used to supplement the lack of fluid in the body in a complex of intensive care measures, as a solvent of other compatible drugs. It can be applied topically for lavage of the wounds, nasal mucosa, as well as catheters and systems for transfusion.
127	NEURO-NORM, 60 caps.	Piracetam Cinnarizine	<i>Rx</i>	N06B X53	Psychostimulant and nootropic. It is prescribed for adults with symptoms of cerebrovascular origin, including impaired memory and thinking functions, reduced concentration, mood disturbances (irritability), labyrinth disorder symptoms, including dizziness, tinnitus, nystagmus, nausea, vomiting; in Meniere's syndrome, for prevention of motion disease, migraine. It is used to treat dyslexia in children aged over 8 years in combination with other appropriate methods.
128	NEOPHYLLINE 50 x 100 mg prolonged release tabs.	Theophylline	<i>Rx</i>	R03D A04	Bronchodilator. It is used for treatment and prevention of bronchospasm associated with bronchial asthma and chronic obstructive pulmonary diseases in adults and children aged over 12 years.
129	NEOPHYLLINE 50 x 300 mg prolonged release tabs.	Theophylline	<i>Rx</i>	R03D A04	

130	NICOTINIC ACID, solution for injection 10 mg/ml, 10 x 1 ml	Nicotinic Acid	<i>Rx</i>	C04A C01	Peripheral vasodilator. It is used to treat pellagra (vitamin PP deficiency), ischemic cerebral circulation disorder, vascular spasm of limbs (obliterating endarteritis, Raynaud's disease), renal vasospasm, poor healing wounds, ulcers, complications of diabetes (diabetic neuropathy, microangiopathy), hyperacid gastritis, enterocolitis, colitis, liver disease, neuritis of the facial nerve, intoxications of different genesis in adults and children aged over 15 years.
131	NIMESULIDE- DARNITSA, 30 x 100 mg tabs.	Nimesulide	<i>Rx</i>	M01A X01	Nonsteroidal anti-inflammatory and antirheumatic drug. It is used to treat acute pain, primary dysmenorrhoea in adults and children aged over 12 years.
132	NIFEDIPINE- DARNITSA, 50 x 0.01 mg film- coated tabs.	Nifedipine	<i>Rx</i>	C08C A05	Calcium channel blockers. It has antianginal and antihypertensive effect. The drug is used for arterial hypertension, ischemic heart disease.
133	NOVOCAINAMIDE- DARNITSA, solution for injection 100 mg/ml, 10 x 5 ml	Procainamide	<i>Rx</i>	C01B A02	Class IA antiarrhythmic drug. It is used in ventricular arrhythmias (extrasystole, tachycardia).
134	NOVOCAIN- DARNITSA, solution for injection 5 mg/ml, 10 x 2 ml amps.	Procaine	<i>Rx</i>	N01B A02	Local anaesthetic. It is prescribed for adults and for local and infiltration anaesthesia, medical block.
135	NOVOCAIN- DARNITSA, solution for injection 5 mg/ml, 10 x 5 ml amps.	Procaine	<i>Rx</i>	N01B A02	
136	NOVOCAIN- DARNITSA, solution for injection 20 mg/ml, 10 x 2 ml amps.	Procaine	<i>Rx</i>	N01B A02	Local anaesthetic. It is prescribed for adults for conduction, epidural and spinal anaesthesia in surgery, urology, ophthalmology, dentistry, otolaryngology, for block of peripheral nerves and nerve plexus.
137	OXOLIN-DARNITSA, 2.5 mg/g ointment 10 g tube	Oxolinum	<i>OTC</i>	S01A D10	Antiviral drug for topical application. It is used for prevention of influenza, treatment of viral rhinitis.

138	OMEPRAZOLE-DARNITSA, 10 x 20 mg caps. in blister pack	Omeprazole	<i>Rx</i>	A02B C01	Drug for treatment of peptic ulcers and gastroesophageal reflux disease. It is used to treat benign gastric and duodenal ulcers, eradication of <i>Helicobacter pylori</i> (in combined therapy with antibacterial drugs), in gastroesophageal reflux disease, to prevent aspiration of acidic gastric contents, Zollinger-Ellison syndrome; reduction of acid-related dyspepsia symptoms.
139	OFLOCAIN-DARNITSA, 0.25 % ointment 1000 g jar	Lidocaine Hydrochloride Ofloxacin	<i>Rx</i>	D06B X50	Antimicrobial drug. It is used to treat complicated infected wounds in the 1st phase of wound healing process of various localization and origin, for prevention and treatment of festering of burn wounds, pyo-inflammatory diseases of skin caused by ofloxacin-sensitive bacterial microflora.
140	OFLOCAIN-DARNITSA, ointment 15 g tube	Lidocaine Hydrochloride Ofloxacin	<i>Rx</i>	D06B X50	
141	OFLOXACIN-DARNITSA, 10 x 200 mg tabs.	Ofloxacin	<i>Rx</i>	J01M A01	Fluoroquinolone antibiotic. The drug is prescribed for adults to treat infectious and inflammatory diseases caused by bacteria susceptible to ofloxacin: - acute and chronic respiratory tract infections; - acute and chronic upper and lower urinary tract infections; - acute and chronic skin and soft tissue infections; - uncomplicated urethral or cervical gonorrhoea; - urethritis and cervicitis of non-gonococcal aetiology.
142	PANTHESTIN-DARNITSA®, gel 15 g tube	Dexpanthenol (D-Panthenol) Myramistin	<i>OTC</i>	D03A X03	The drug promotes wound healing. It is used to treat wounds of different localization and genesis, trophic ulcers and decubitus in the regeneration phase (phase II of wound healing), to prepare the granulating burn wounds to autodermoplasty, to improve engraftment of skin transplants after plastic surgery, for treatment of sunburns, for prevention of radiation lesions of the mucous membranes, skin and surrounding tissues, treatment of radiation ulcers in the phase of regeneration, treatment of periodontal diseases, diseases of the oral mucosa, treatment of cervical erosion, erosive colpitis and mucosal lesions of the vagina, skin infections, minor abrasions, cuts, scratches, burns and prevention of infection in home traumatism. In paediatric patients the drug is used in diaper rash and diaper dermatitis.
143	PANTHESTIN-DARNITSA®, gel 30 g tube	Dexpanthenol (D-Panthenol) Myramistin	<i>OTC</i>	D03A X03	
144	PAPAVERIN-DARNITSA, solution for injection 20 mg/ml, 10 x 2 ml	Papaverine Hydrochloride	<i>Rx</i>	A03A D01	Myotropic antispasmodic. It used in spasms of smooth muscles of the abdomen (pylorospasm, spastic colitis, cholecystitis, attacks of gallstone disease), urinary tract, renal colic, cerebral spasms, coronary spasms (related, for example, to acute myocardial infarction, angina pectoris), peripheral vasospasm (endarteritis). The drug is prescribed for adults and children aged over 1 year.
145	PAPAZOLUM-DARNITSA, 10 tabs.	Bendazolu Hydrochloride Papaverine Hydrochloride	<i>OTC</i>	C04A X51	Myotropic antispasmodic. It is used in spasms of smooth muscles of internal organs (pylorospasm, spastic colitis, cholecystitis, peptic ulcer and duodenal ulcer, intestinal colic, renal colic, urinary tract spasm), transient cerebral vasospasm, moderate peripheral vasospasm, in the combined therapy of arterial hypertension.

146	PARACETAMOL-DARNITSA, 10 x 200 mg tabs.	Paracetamol	<i>OTC</i>	N02B E01	Analgesic and antipyretic drug. It is used to treat mild to moderate pain of different genesis, relief of symptoms of colds and influenza (fever, aches) in adults and children aged over 3 years.
147	PARACETAMOL-DARNITSA, 10 x 500 mg tabs.	Paracetamol	<i>OTC</i>	N02B E01	
148	PENTOXIFYLLINE-DARNITSA, solution for injection 20 mg/ml, 10 x 5 ml	Pentoxifylline	<i>Rx</i>	C04A D03	Peripheral vasodilator. It is used in atherosclerotic encephalopathy, cerebral ischemic stroke, discirculatory encephalopathy, peripheral circulation disorders, trophic disorders in the tissues associated vein injuries or microcirculation disorders, obliterating endarteritis, Raynaud's disease, eye circulatory disorders (acute, subacute, chronic insufficiency of blood circulation in the retina and choroid), dysfunction of the inner ear of vascular genesis, accompanied by hearing loss.
149	PENTOXIFYLLINE-DARNITSA, 20 x 200 mg tabs.	Pentoxifylline	<i>Rx</i>	C04A D03	
150	PICONORM, oral drops, 7.5 mg/ml solution, 1 x 10 ml	Sodium Picosulfate	<i>OTC</i>	A06A B08	Triarylmethane laxative which after metabolism in the large intestine inhibits absorption of water and thus increases the excretion of water and electrolytes. This results in softening of stool consistency and rapid bowel evacuation, increasing the volume of stool and stimulating peristalsis. It is used for short-term use in constipations or conditions requiring to facilitate defecation. The drug is contraindicated in children aged below 4 years. The drug is used only on prescription in children aged over 4 years.
151	PIPERAZINE ADIPINATE-DARNITSA, 10 x 200 mg tabs.	Piperazini Adipate	<i>Rx</i>	P02C B01	Anthelmintic drug. It is used in ascariasis, enterobiasis in adults and children aged over 4 years.
152	PIRACETAM-DARNITSA, solution for injection 200 mg/ml, 10 x 5 ml	Piracetam	<i>Rx</i>	N06B X03	Psychostimulant and nootropic. It is prescribed for adults for: ^ - symptomatic treatment of psycho-organic syndrome, accompanied by loss of memory, dizziness, decreased concentration; ^ - treatment of vertigo and related disorders of balance, with the exception of dizziness and vasomotor and mental origin; ^ - treatment of cortical myoclonus as monotherapy or in combined therapy; ^ - treatment of chronic alcoholism and alcohol withdrawal syndrome; ^ In children aged below 8 years it is used to treat dyslexia in combination with other appropriate methods, including speech therapy.
153	PIRACETAM-DARNITSA, 30 x 400 mg tabs.	Piracetam	<i>Rx</i>	N06B X03	Psychostimulant and nootropic. Indications in adults: - symptomatic treatment of psycho-organic syndrome, accompanied by loss of memory,

					<p>decreased concentration and general activity;</p> <ul style="list-style-type: none"> - treatment of vertigo and related disorders of balance, with the exception of dizziness and vasomotor and mental origin; - treatment of cortical myoclonus as monotherapy or in combined therapy; - in combined treatment of sickle cell anaemia to reduce the manifestations of the vaso-occlusive crisis. <p>Indications in children: treatment of dyslexia in children aged over 8 years in combination with other appropriate methods, including speech therapy; treatment of cortical myoclonus as monotherapy or in combined therapy; in combined treatment of sickle cell anaemia in children aged over 3 years to reduce the manifestations of the vaso-occlusive crisis.</p>
154	PIRACETAM-DARNITSA, 60 x 200 mg film-coated tabs.	Piracetam	<i>OTC</i>	N06B X03	<p>Psychostimulant and nootropic.</p> <p>It is prescribed for adults for:</p> <ul style="list-style-type: none"> - symptomatic treatment of psycho-organic syndrome, accompanied by loss of memory, dizziness, decreased concentration; - treatment of vertigo and related disorders of balance, with the exception of dizziness and vasomotor and mental origin; - treatment of cortical myoclonus as monotherapy or in combined therapy; - treatment of chronic alcoholism and alcohol withdrawal syndrome; <p>In children aged below 8 years it is used to treat dyslexia in combination with other appropriate methods, including speech therapy.</p>
155	PYRIDOXINE-DARNITSA (VITAMIN B6-DARNITSA), solution for injection 50 mg/ml, 10 x 1 ml amps.	Pyridoxine	<i>Rx</i>	A11H A02	<p>Vitamin. It plays an important role in the metabolism, necessary for normal functioning of the central and peripheral nervous system.</p> <p>Indications: Hypo- and vitamin B6 deficiency, combined treatment of toxemia of pregnancy, atherosclerosis, anaemia, leucopenia, nervous system diseases, involutional depression, seborrheic and non-seborrheic dermatitis, herpes zoster, neurodermatitis, psoriasis, exudative diathesis, elimination of binge-drinking and hangover syndrome.</p>
156	PLATYPHYLLINE-DARNITSA, solution for injection 2 mg/ml, 10 x 1 ml	Platyphylline Hydrotartrate	<i>Rx</i>	A03A X12	<p>Natural alkaloid that has M-anticholinergic action.</p> <p>It is used in combined therapy of gastroduodenitis, functional dyspepsia, pylorospasms, cholecystitis, cholelithiasis, intestinal colic, renal colic, biliary colic.</p> <p>The drug is prescribed for treatment of bronchial asthma to prevent bronchospasm, bronchorrhoea; in algodysmenorrhoea; spasm of cerebral arteries; angiotrophoneurosis.</p>
157	PREDNISOLONE-DARNITSA, 40 x 5 mg tabs.	Prednisolone	<i>Rx</i>	H02A B06	<p>The drug has anti-inflammatory, antiallergic, desensitizing, antishock and immunosuppressive effect.</p> <p>Indications for use:</p> <ul style="list-style-type: none"> -rheumatic fever, rheumatic carditis, chorea; - systemic connective tissue disease; -multiple sclerosis; -acute and chronic inflammatory diseases of the joints; -bronchial asthma, asthmatic status; -interstitial lung disease, lung cancer (in combination with cytotoxic), berylliosis, aspiration pneumonia (in combination with specific therapy), eosinophilic Loeffler's pneumonia, tuberculosis (pulmonary tuberculosis, tuberculous meningitis) - combined with specific therapy ;

					- primary and secondary adrenal insufficiency (including conditions after removal of the adrenal glands), congenital adrenal hyperplasia, adrenogenital syndrome, subacute thyroiditis; - acute and chronic allergic diseases ; -autoimmune diseases; - see detailed indications in the instruction for medical use
158	PREDNISOLONE-DARNITSA, solution for injection 30 mg, 3 x 1 ml	Prednisolone	<i>Rx</i>	H02A B06	The drug has anti-inflammatory, antiallergic, desensitizing, antishock and immunosuppressive effect. Indications for use: Inflammatory joint disease (rheumatoid arthritis etc.) acute rheumatism; Addison's disease; acute failure of the adrenal cortex ; adrenogenital syndrome; bronchial asthma; asthmatic status; acute and chronic diseases requiring rapid increase in concentrations of glucocorticoids in the body (increased sensitivity to drugs, serum disease, urticaria, allergic rhinitis, hypersensitivity to chemicals, angioedema); hepatitis; hepatic coma; hypoglycaemic conditions; lipoid nephrosis; blood system diseases (agranulocytosis, various forms of leukaemia, thrombocytopenic purpura, haemolytic anaemia); Hodgkin's disease; see more details on application in instructions.
159	PREDNICARBUM-DARNITSA, ointment 15 g tube	Prednisolone Urea	<i>OTC</i>	D07X A02	Combined drug for topical use with anti-inflammatory, antiexudative, antiallergic, antipruritic, keratoplastic, keratolytic, antimicrobial action. It is used for treatment of psoriasis; atopic dermatitis; various forms of eczema (including lichenification); dyshidrosis; seborrhoeic dermatitis; ichthyosis; ichthyosiform dermatoses, accompanied by dryness and hyperkeratosis.
160	PROSERIN-DARNITSA, solution for injection 0.5 mg/ml, 10 x 1 ml	Neostigmine Bromide	<i>Rx</i>	N07A A01	Synthetic reversible cholinesterase inhibitor. Indications for use: -myasthenia , acute myasthenic crisis ; -motor violations after brain injury; -paralysis; -recovery period after meningitis, polio; encephalitis ; -neuritis, optic atrophy ; -intestinal atony, bladder atony; - elimination of residual effects after blockade of neuromuscular transmission with nondepolarizing muscle relaxant.
161	RANITIDINE-DARNITSA, 10 x 150 mg film-coated tabs.	Ranitidine	<i>Rx</i>	A02B A02	Antiulcer drug, Histamine H2-receptor antagonist. The drug reduces basal and stimulated secretion of hydrochloric acid, reducing the amount of gastric juices caused by stimulation of baroreceptors (stretching of the stomach), nutritional load, action of hormones and biogenic stimulators (gastrin, histamine, pentagastrin, caffeine).
162	RANITIDINE-DARNITSA, 20 x 150 mg film-coated tabs.	Ranitidine	<i>Rx</i>	A02B A02	Indications for use: - peptic ulcer of the stomach and duodenum, not associated with Helicobacter pylori (in exacerbation phase), including ulcers, associated with non-steroidal anti-inflammatory drug administration (NSAIDs) ; - functional dyspepsia ; - chronic gastritis with increased stomach acid-forming function in the acute stage ; - gastroesophageal reflux disease (to relieve symptoms) or reflux esophagitis.

163	REMISID, 10 mg gel 30 g tube	Nimesulide	<i>Rx</i>	M01A X17	Nonsteroidal anti-inflammatory drug. In topical application causes decrease or disappearance of pain in the area of application, including joint pain, at rest and during movement, reduces morning stiffness and swelling of joints, increases range of motion. It is used for the topical treatment of pathological conditions of the musculoskeletal system (characterized by pain, inflammation and stiffness of movement), such as osteoarthritis, periarthritis, post-traumatic tendinitis, tendosynovitis, muscle stretching, severe physical load on the joints.
164	RIBOXINUM-DARNITSA, solution for injection 20 mg/ml, 10 x 10 ml	Inosine	<i>Rx</i>	C01E B14	Anabolic drug that has antihypoxic and antiarrhythmic action. The drug normalizes myocardial contractile activity and stimulates complete myocardial relaxation in diastole through the ability to bind calcium ions that penetrated cells during stimulation, activates tissue regeneration. Indications for use: -combined therapy of coronary heart disease: -arrhythmia; -intoxication with cardiac glycosides; -treatment of cardiomyopathies of various genesis, yocardiodystrophy, myocarditis; liver disease; - prevention of leucopenia under radiation therapy; to improve visual function in open-angle glaucoma with normalized intraocular pressure.
165	RIBOXINUM-DARNITSA, solution for injection 20 mg/ml, 10 x 5 ml	Inosine	<i>Rx</i>	C01E B14	Anabolic drug that has antihypoxic and antiarrhythmic action. Indications for use: - combined therapy of coronary heart disease (condition after myocardial infarction, angina pectoris); -arrhythmia; -intoxication with cardiac glycosides; -treatment of cardiomyopathies of various genesis, myocardiodystrophy, myocarditis; -liver diseases (hepatitis, liver cirrhosis, hepatosteatosi); -urocoproporphyria.
166	RIBOXINUM-DARNITSA, 50 x 200 mg film-coated tabs.	Inosine	<i>Rx</i>	C01E B14	Anabolic drug that has antihypoxic and antiarrhythmic action. Indications for use: - combined therapy of coronary heart disease (condition after myocardial infarction, angina pectoris); -arrhythmia; -intoxication with cardiac glycosides; -treatment of cardiomyopathies of various genesis, myocardiodystrophy, myocarditis; -liver diseases (hepatitis, liver cirrhosis, hepatosteatosi); -urocoproporphyria.
167	RIMANTADINE-DARNITSA, 20 x 50 mg tabs.	Rimantadine	<i>OTC</i>	J05A C02	Antiviral drug. It is used for prevention and treatment of influenza caused by type A virus, prevention of tick-borne encephalitis of virus aetiology in adults.
168	RINAZAL®, nasal drops, 1 mg/ml, 10 ml vial	Xylometazoline	<i>OTC</i>	R01A A07	Drug for local use in diseases of the nasal cavity. It is used for the symptomatic treatment of nasal congestion in colds, hay fever, allergic rhinitis, sinusitis ; to facilitate the secretion outflow in diseases of the sinuses; as adjunctive therapy of otitis media (to reduce swelling of nasal mucosa); to facilitate rhinoscopy.
169	ROXILID®, 10 x 150 mg film-coated tabs.	Roxithromycin	<i>Rx</i>	J01F A06	Semisynthetic second generation macrolide antibiotic. The drug is prescribed to treat infections caused by microorganisms susceptible to roxithromycin: – ENT organs (tonsillitis, laryngitis, pharyngitis, sinusitis, otitis media); - respiratory tract infections (pneumonia, acute and chronic bronchitis); - skin and soft tissue (common acne, furunculus, carbuncle, pyoderma); - urogenital infections (except gonorrhoea, including urethritis, cervicovaginitis).

170	SEPTEFRILUM-DARNITSA, 10 x 0.2 mg tabs.	Decametoxine	<i>OTC</i>	R02A A20	Drugs for treatment of throat diseases. It has a wide spectrum of antimicrobial action. It is used for local treatment in combined therapy of diseases of the oral cavity, pharynx, larynx (pharyngitis, laryngitis, tonsillitis, gingivitis, periodontitis, stomatitis, oral candidiasis, initial tonsillitis), and for prevention of infectious complications after surgical interventions in the oral cavity, pharynx, larynx.
171	SEPTEFRILUM-DARNITSA, 20 x 0.2 mg tabs.	Decametoxine	<i>OTC</i>	R02A A20	
172	SPIRONOLACTONE-DARNITSA, 30 x 25 mg tabs.	Spirolactone	<i>Rx</i>	C03D A01	Potassium-sparing diuretic. Aldosterone antagonist. It is prescribed to treat the following conditions: - primary hyperaldosteronism; - congestive heart failure after ineffectiveness or intolerance to other diuretics, in case of required increase of their efficiency ; - essential arterial hypertension, mainly in hypokalaemia, usually in combination with other antihypertensive drugs; - liver cirrhosis accompanied by oedema and/or ascites; - swelling caused by nephrotic syndrome; -hypokalaemia - in case of inability to apply other therapy. It is used for prevention of hypokalaemia in patients receiving cardiac glycosides, if other methods of treatment are considered as inappropriate or irrelevant.
173	STREPTOCIDE-DARNITSA, 10 X 300 mg tabs.	Sulfanilamide	<i>OTC</i>	J01E B06	The drug has bacteriostatic effect on streptococci, meningococci , gonococci, Escherichia coli, toxoplasmosis and malaria pathogens. It is used to treat infectious and inflammatory diseases caused by microorganisms susceptible to the drug: infections of the skin and mucous membranes (wounds, ulcers, decubitus), enterocolitis, pyelitis, cystitis.
174	STROPHANTIN-DARNITSA, solution for injection 0.25 mg/ml, 10 x 1 ml	Strophanthin	<i>Rx</i>	C01A C01	The drug has cardiotoxic action, increases the force and velocity of myocardial contractions (positive inotropic effect), reduces heart rate (negative chronotropic effect), reduces atrioventricular conduction (negative dromotropic effect). In heart failure it increases cardiac stroke volume and minute volume, improves ventricular emptying, leading to improvement in blood flow. It is used in acute heart failure, chronic heart failure stage IIb-III (III-IV stage according to NYHA classification), supraventricular tachycardia, atrial fibrillation.
175	SULFADIMETHOXINE-DARNITSA, 10 x 500 mg tabs.	Sulfadimethoxine	<i>Rx</i>	J01E D01	Antimicrobial bacteriostatic drug, long-acting sulphonamide. It is used to treat infectious and inflammatory diseases caused by microflora sensitive to the drug: tonsillitis, bronchitis, pneumonia, sinusitis, otitis, dysentery, inflammatory diseases of the biliary and urinary tract, gonorrhoea, erysipelas, pyoderma, meningitis, infection of wounds, trachoma, toxoplasmosis, shigellosis, resistant forms of malaria (in combination with antimalarial drugs).
176	SULFOCAMPHOCAIN-DARNITSA, solution for injection 100 mg/ml, 10 x 2 ml	Novocaine (Procaine Hydrochloride) Sulfokamforna Acid	<i>Rx</i>	C01E B02	The drug tones the respiratory and vasomotor centres, increases metabolism in the heart muscle, increasing its sensitivity to influence of sympathetic nerves. [It is used to treat acute and chronic respiratory failure; acute heart failure in geriatrics; respiratory depression in pneumonia and other infectious diseases; cardiogenic and anaphylactic shock; alcohol intoxication, mild poisoning with hypnotics.

177	TAUFONE-DARNITSA, 40 mg/ml eye drops, 10 ml	Taurine	<i>Rx</i>	S01X A21*	The drug stimulates reparative and regenerative processes in dystrophic eye diseases and/or pathological processes, accompanied by severe disturbance of the metabolism of eye tissues, including eye injuries. Promotes normalisation of the function of cell membranes, optimization of energy metabolism, support of electrolyte sustainability of the cells cytoplasm, inhibition of synaptic transmission. Indications: -dystrophic lesions of the cornea and retina, including hereditary retinal abiotrophy; -cataract (age-related, diabetic, traumatic, radiation); -as a drug for stimulating reparative processes in cornea injuries; -open-angle glaucoma (as additional drug).
178	TIMOLOL-DARNITSA, eye drops, 5 mg/ml, 5 ml vial	Timolole	<i>Rx</i>	S01E D01	Timolol is β_1 - and β_2 -adrenergic receptor blocker. In instillation in the eye it reduces intraocular pressure mainly by reducing the production of intraocular liquid. It does not affect accommodation, refraction and pupil size. It is used in eye hypertension, chronic open-angle glaucoma, in some cases of secondary glaucoma.
179	THIAMINE CHLORIDE- DARNITSA (VITAMIN B1- DARNITSA), solution for injection 50 mg/ml, 10 x 1 ml	Thiamine	<i>Rx</i>	A11D A01	Synthetic vitamin B1 supplement, belongs to water-soluble vitamins. Vitamin B1 plays an important role in metabolism and neuro-reflex regulation. It is used for treatment of hypo- and vitamin B1 deficiency, neuritis, polyneuritis, radiculitis, neuralgia, peripheral paralysis, encephalopathy, neurasthenia, gastric ulcer and duodenal ulcer, intestinal atony, chronic liver disease, myocardial dystrophy, endarteritis, dermatitis of neurogenic origin, pyoderma, eczema, psoriasis.
180	TIVOMAX- DARNITSA, solution for infusion 42 mg/ml, 1 x 100 ml	Arginine Hydrochloride	<i>Rx</i>	B05X B01	The active ingredient of the drug - arginine - is an active cellular regulator of many vital body functions, has important protective effects for the body in critical state. The drug has antihypoxic, membrane-stabilizing, cell protective, antioxidant, antiradical, detoxification effects. It is used in atherosclerosis, angiopathy, ischemic heart disease, congestive heart failure, arterial hypertension, hypercholesterolemia, lung diseases and hypoxic conditions, in hepatitis, hyperammonemia, asthenic conditions, thymus function reduction, delayed foetal development, in eclampsia.
181	TONORMA®- DARNITSA, 10 film-coated tabs.	Atenolol, Chlorthalidone , Nifedipine	<i>Rx</i>	C07F B0	Combined antihypertensive drug containing atenolol, nifedipine and chlorthalidone. The drug has antihypertensive, antianginal and antiarrhythmic effects.
182	TONORMA®- DARNITSA, 30 film-coated tabs.	Atenolol, Chlorthalidone , Nifedipine	<i>Rx</i>	C07F B0	
183	TRIMISTIN®- DARNITSA, ointment 14 g tube	Triamcinolone Acetonide Micronized Myramistin	<i>OTC</i>	D07B B03	Combined drug for local use. The drug has a pronounced antibacterial, anti-inflammatory, anti-allergic and antiexudative action. It is used to treat inflammatory skin diseases, particularly complicated by bacterial or fungal infections: eczema, atopic dermatitis, neurodermatitis, psoriasis in advanced stage and in exudative form,

					parapsoriasis, pemphigus, contact dermatitis, seborrheic dermatitis, Duhring dermatitis, discoid lupus erythematosus, photodermatosis, skin lymphoma.
184	TRIFTAZIN-DARNITSA®, solution for injection 2 mg/ml, 10 x 1 ml	Trifluoperazine	<i>Rx</i>	N05A B06	Antipsychotic drug (neuroleptic) iperazine phenothiazine derivative. The drug has antipsychotic, sedative, antiemetic, cataleptic, hypotensive, hypothermic and mild anticholinergic effect, also against hiccups. It is used to treat psychotic disorders, including schizophrenia.
185	TROXERUTIN-DARNITSA®, 20 mg/g gel 30 g tube	Troxerutine	<i>OTC</i>	C05C A04	The action of the drug is aimed at pain reduction, improvement of various pathological disorders associated with venous insufficiency. It is indicated for the symptomatic treatment of: venous insufficiency and pre-varicose and varicose syndrome, superficial thrombophlebitis, phlebitis and post-phlebitis conditions; for the combined therapy of haemorrhoidal disease; swelling and pain in injuries and varicose veins; convulsive contractures of calf muscles.
186	T-TRIOMAX, solution for injection 25 mg/ml, 10 x 4 ml	Morfolinijeva Acidum Salis Tiazotnoyi	<i>Rx</i>	C01E B	Cardiovascular drug, it has anti-ischemic, membrane stabilizing, antioxidant and immunomodulating effect. It is used in combined therapy of coronary heart disease, chronic hepatitis, alcoholic hepatitis, liver fibrosis and cirrhosis, as an additional drug in treatment of cardiac arrhythmias.
187	T-TRIOMAX, solution for injection 25 mg/ml, 10 x 2 ml	Morfolinijeva Acidum Salis Tiazotnoyi	<i>Rx</i>	C01E B	Cardiovascular drug, it has anti-ischemic, membrane stabilizing, antioxidant and immunomodulating effect. It is used in combined therapy of coronary heart disease, chronic hepatitis, alcoholic hepatitis, liver fibrosis and cirrhosis, as an additional drug in treatment of cardiac arrhythmias.
188	URSOCHOL®, 100 x 250 mg caps.	Ursodeoxycholic Acid	<i>Rx</i>	A05A A02	After oral administration ursodeoxycholic acid decreases the cholesterol saturation of bile, inhibiting its absorption in the intestine and reducing the secretion of cholesterol in bile. Indications: - dissolution of roentgen-negative cholesterol gallstones no larger than 15 mm in diameter in patients with functioning gallbladder, despite the presence of gallstone(-s); - treatment of gastritis with bile reflux; - symptomatic treatment of primary biliary cirrhosis (PBC) in the absence of decompensated liver cirrhosis.
189	URSOCHOL®, 50 x 250 mg caps.	Ursodeoxycholic Acid	<i>Rx</i>	A05A A02	After oral administration ursodeoxycholic acid decreases the cholesterol saturation of bile, inhibiting its absorption in the intestine and reducing the secretion of cholesterol in bile. Indications: - dissolution of roentgen-negative cholesterol gallstones no larger than 15 mm in diameter in patients with functioning gallbladder, despite the presence of gallstone(-s); - treatment of gastritis with bile reflux; - symptomatic treatment of primary biliary cirrhosis (PBC) in the absence of decompensated liver cirrhosis.
190	FAMOTIDINE-DARNITSA, 20 x 20 mg film-coated tabs.	Famotidin	<i>Rx</i>	A02B A03	Famotidine is a histamine H2-receptor blocker of stomach wall, so it reduces the secretion of gastric juice. Under the influence of the drug both the concentration and amount of gastric juice is reduced. It is used in benign gastric ulcers, peptic duodenal ulcer (treatment and prevention of relapses), in hypersecretory conditions, such as Zollinger-Ellison syndrome. It is used for treatment of gastroesophageal reflux disease (reflux esophagitis), for prevention of symptoms and erosions or ulceration associated with gastroesophageal reflux disease.

191	F-GEL®, 25 mg/g gel 30 g tube	Ketoprofen	<i>Rx</i>	MO2A A10	Nonsteroidal anti-inflammatory drug for topical application. It is used in muscle and joint pain caused by injuries or damages; for treatment of tendovaginitis.
192	FLUCONAZOLE-DARNITSA, 10 x 50 mg caps.	Fluconazole	<i>Rx</i>	J02A C01	Antifungal agent. It is used for treatment of cryptococcal meningitis; coccidioidomycosis; invasive candidiasis; mucosal candidiasis, chronic candidiasis of the skin and mucous membranes; chronic atrophic candidiasis after failure of local dental hygiene products; vaginal candidiasis, acute or recurrent, when local therapy is not appropriate; Candida balanitis; dermatomycoses, einea cruris, tinea versicolor, Candida infections of the skin; dermatophytic onychomycosis.
193	FLUCONAZOLE-DARNITSA, 10 x 100 mg caps.	Fluconazole	<i>Rx</i>	J02A C01	
194	FLUCONAZOLE-DARNITSA, 1 x 150 mg caps.	Fluconazole	<i>OTC</i>	J02A C01	
195	FLUCONAZOLE-DARNITSA, 2 x 150 mg caps.	Fluconazole	<i>OTC</i>	J02A C01	
196	FLUCONAZOLE-DARNITSA, 3 x 150 mg caps.	Fluconazole	<i>OTC</i>	J02A C01	
197	FLUCONAZOLE-DARNITSA, solution for infusion 2 mg/ml, 1 x 100 ml vial	Fluconazole	<i>Rx</i>	J02A C01	Antifungal drug for system use. It is prescribed for adults and children from birth. It is indicated for treatment of candidiasis, coccidioidomycosis, cryptococcal meningitis and prevention of this disease in groups of increased risk of disease.
198	FLUCAR-DARNITSA, 1 mg/g cream 15 g tube	Fluocinoloni Acetonide	<i>Rx</i>	D07A C04	Corticosteroids for use in dermatology. The drug is used for short-term local treatment of acute and severe non-infectious skin diseases (dry forms), scalp psoriasis, seborrhoeic dermatitis, lichen planus.

199	FOCORT®-DARNITSA, 1 mg/g cream 15 g tube	Triamcinolone	<i>Rx</i>	D07A B09	The drug has anti-inflammatory, anti-allergic and antipruritic effect. It is used for treatment of all types of acute and chronic eczema regardless of location; neurodermatitis; contact dermatitis, regardless of aetiology ; red lichen planus; vulgar psoriasis; exfoliative erythroderma; Leiner's disease; otitis externa uncomplicated by infection; lupus erythematosus.
200	PHTHALAZOL-DARNITSA, 10 x 500 mg tabs.	Phthalylsulfathiazole	<i>OTC</i>	A07A B02	Antimicrobial drug. It is used to treat acute dysentery (shigellosis), chronic dysentery in the acute phase, colitis, enterocolitis, gastroenteritis; for prevention of infectious complications during intestinal surgeries.
201	FUROSEMIDE-DARNITSA, solution for injection 10 mg/ml, 10 x 2 ml	Furosemide	<i>Rx</i>	C03C A01	Furosemide is a fast acting loop diuretic with relatively strong short-term diuretic effect. It is used for treatment of: - swelling in chronic congestive heart failure (if required treatment with diuretics); - swelling in acute congestive heart failure; - swelling in chronic renal failure; - acute renal failure, including acute renal failure in pregnant women or during childbirth; - swelling in liver diseases (if required, to supplement treatment with aldosterone antagonists); - hypertensive crisis (as a supporting drug).
202	FUROSEMIDE-DARNITSA, 50 x 40 mg tabs.	Furosemide	<i>Rx</i>	C03C A01	Furosemide is a fast acting loop diuretic with relatively strong short-term diuretic effect. It is used for treatment of: - swelling in chronic congestive heart failure (if required treatment with diuretics); - swelling in nephrotic syndrome (if required treatment with diuretics); - swelling in chronic renal failure; - acute renal failure, including acute renal failure in pregnancy or during childbirth; - swelling in liver diseases (if required, to supplement treatment with aldosterone antagonists); - arterial hypertension.
203	CITIMAX-DARNITSA, solution for injection 250 mg/ml, 5 x 4 ml	Citicolinum	<i>Rx</i>	N06BX06	Psycho-stimulant and nootropic drug Indications for use: - acute phase of cerebral circulation disorders and treatment of their consequences ; - traumatic brain injury and its consequences ; - neurological disorders (cognitive, sensitive, motor) caused by cerebral pathology of degenerative and vascular origin.
204	CITIMAX-DARNITSA, solution for injection 250 mg/ml, 10 x 4 ml	Citicolinum	<i>Rx</i>	N06BX06	
205	CEFAZOLIN-DARNITSA, powder for solution for injection, 5 x 0.5 g vial	Cefazolin	<i>Rx</i>	J0D A04	Antimicrobial drug for systemic use. Semisynthetic first generation cephalosporin antibiotic for parenteral administration, broad-spectrum action, active against many gram-positive and gram-negative microorganisms. It is used to treat infections caused by organisms susceptible to cefazolin; - respiratory tract infections; - genitourinary system infections; -

206	CEFAZOLIN-DARNITSA, powder for solution for injection, 5 x 1.0 g vial	Cefazolin	<i>Rx</i>	J0D A04	skin and soft tissue infections; - infections of bones and joints; - sepsis; - endocarditis; - biliary tract infections. Prevention of surgical infections.
207	CEFOKTAM, powder for solution for injection, 1 x 0.75 g vial	Cefuroxim	<i>Rx</i>	J01D C02	Antimicrobial drug for systemic use. Second generation cephalosporin antibiotic for parenteral use. It has bactericidal action, violates the synthesis of cell walls of microorganisms. It has a wide range of action. The drug is used to treat infectious and inflammatory diseases caused by pathogen susceptible to drug, including upper and lower respiratory tract infections - acute and chronic bronchitis, bacterial pneumonia, lung abscess, postoperative infections of the chest cavity; infections of ear, throat, nose - sinusitis, tonsillitis, pharyngitis and otitis media; urinary tract infections; inflammatory diseases of the pelvic organs (in obstetrics and gynaecology); gonorrhoea; other infections - peritonitis, septicaemia, meningitis; prevention of infections during surgeries on the organs of the chest and abdomen, pelvic organs, during cardiovascular and orthopaedic surgeries.
208	CEFOKTAM, powder for solution for injection, 1 x 1.5 g vial	Cefuroxim	<i>Rx</i>	J01D C02	Antimicrobial drug for systemic use. Third generation cephalosporin antibiotic for parenteral use. It has bactericidal action. It has a wide range of actions. It is used to treat infections caused by microorganisms susceptible to the drug: - upper respiratory tract infection (tonsillitis, otitis); - respiratory tract infections (bronchitis, pneumonia, pleurisy, abscesses); - genitourinary system infections; - septicaemia, bacteraemia; - intra-abdominal infections (including peritonitis); - skin and soft tissue infections; - infections of bones and joints; - meningitis (except listeriosis) and other infections of the central nervous system. Also to prevent infections after surgeries on the digestive tract, urological and obstetrical surgeries.
209	CEFOTAXIME-DARNITSA, powder for solution for injection, 1 x 1.0 g vial	Cefotaxime	<i>Rx</i>	J0D A10	Antimicrobial drug for systemic use. Third generation cephalosporin antibiotic for parenteral use. It has bactericidal action. It has a wide range of actions. It is used to treat infections caused by microorganisms susceptible to the drug: - upper respiratory tract infection (tonsillitis, otitis); - respiratory tract infections (bronchitis, pneumonia, pleurisy, abscesses); - genitourinary system infections; - septicaemia, bacteraemia; - intra-abdominal infections (including peritonitis); - skin and soft tissue infections; - infections of bones and joints; - meningitis (except listeriosis) and other infections of the central nervous system. Also to prevent infections after surgeries on the digestive tract, urological and obstetrical surgeries.
210	CEFOTAXIME-DARNITSA, powder for solution for injection, 5 x 1.0 g vial	Cefotaxime	<i>Rx</i>	J0D A10	Antimicrobial drug for systemic use. Third generation cephalosporin antibiotic for parenteral use. It has bactericidal action. It has a wide range of actions. It is used to treat infections caused by microorganisms susceptible to the drug: - upper respiratory tract infection (tonsillitis, otitis); - respiratory tract infections (bronchitis, pneumonia, pleurisy, abscesses); - genitourinary system infections; - septicaemia, bacteraemia; - intra-abdominal infections (including peritonitis); - skin and soft tissue infections; - infections of bones and joints; - meningitis (except listeriosis) and other infections of the central nervous system. Also to prevent infections after surgeries on the digestive tract, urological and obstetrical surgeries.
211	CEFTRIAXONUM-DARNITSA, powder for solution for injection, 1 x 0.5 g vial	Ceftriaxone	<i>Rx</i>	J0D A13	Antimicrobial drug for systemic use. It is used to treat infections, caused by pathogens sensitive to ceftriaxone: - respiratory tract infections, particularly pneumonia, and infections of ear, throat and nose; - genital infections, including gonorrhoea; - septicaemia; - infections of bones, joints, soft tissues, skin and wound infections;
212	CEFTRIAXONUM-DARNITSA, powder for solution for injection, 5 x 0.5 g vial	Ceftriaxone	<i>Rx</i>	J0D A13	- infections in patients with neutropenia; - meningitis. Also, for preoperative infection prevention during surgeries on the organs of the gastrointestinal tract, biliary tract, urinary tract during gynaecological procedures, but only in cases of known or potential contamination.

213	CEFTRIAXONUM-DARNITSA, powder for solution for injection, 1 x 1.0 g vial	Ceftriaxone	<i>Rx</i>	J0D A13	
214	CEFTRIAXONUM-DARNITSA, powder for solution for injection, 40 x 1.0 g vial	Ceftriaxone	<i>Rx</i>	J0D A13	
215	CEFTRIAXONUM-DARNITSA, powder for solution for injection, 5 x 1.0 g vial	Ceftriaxone	<i>Rx</i>	J0D A13	
216	CITRAMON EXTRA, 10 tabs.	Paracetamol, Caffeine	<i>OTC</i>	N02B A51	Combined drug. It has analgesic, antipyretic and anti-inflammatory effect, helps normalisation of cerebral vascular tone and blood flow acceleration. It is used to treat mild to moderate pain syndrome (migraine, headache, back pain, toothache, neuralgia, myalgia, arthralgia, rheumatic pain, primary dysmenorrhoea, periodic pain in women), as well as antipyretic drug in fever of different aetiology , in catarrhal diseases, influenza and inflammation of the throat.
217	CITRAMON-DARNITSA, 6 tabs.	Acetylsalicylic Acid, Paracetamol, Caffeine	<i>OTC</i>	N02B A51	The drug has analgesic, antipyretic and anti-inflammatory action. It is used as a therapy of mild to moderate pain syndrome: in headache or dental pain, in primary dysmenorrhoea, migraine, arthralgia, neuralgia, diseases accompanied by hyperthermia of various aetiologies (as antipyretic drug).
218	CITROPAK-DARNITSA, 6 tabs.	Acetylsalicylic Acid, Paracetamol, Caffeine	<i>OTC</i>	N02B A51	Combination drug. It is used to treat mild to moderate pain syndrome (headache, migraine, toothache, sore throat, neuralgia , inflammation of the fibrous tissue, myalgia, arthralgia, pain in dislocations and sprains, rheumatic pain, sciatica, lumbago, primary dysmenorrhoea), feverish conditions of different origin.
219	CYANOCOBALAMIN-DARNITSA (VITAMIN B12-DARNITSA), solution for injection 0.2 mg/ml, 10 x 1 ml amps.	Cyanocobalamin	<i>Rx</i>	B03B A01	Vitamin B12. It has metabolic, hematopoietic effect. It is used in chronic anaemia, chronic hepatitis, liver cirrhosis, liver failure, polyneuritis, radiculitis, neuralgia, skin diseases, diseases of the stomach and intestine, accompanied by malabsorption of vitamin B12, in acute radiation disease.
220	CYANOCOBALAMIN-DARNITSA (VITAMIN B12-DARNITSA), solution for injection 0.5 mg/ml, 10 x 1 ml amps.	Cyanocobalamin	<i>Rx</i>	B03B A01	

